

ON AUTOMORPHISMS OF C^* -ALGEBRAS WHOSE VOICULESCU ENTROPY IS GENUINELY NONCOMMUTATIVE

ADAM SKALSKI

ABSTRACT. We use the results of Neshveyev and Størmer to show that for a generic shift on a C^* -algebra associated to a bitstream the Voiculescu topological entropy is strictly larger than the supremum of topological entropies of its classical subsystems.

Let A be an (exact) C^* -algebra, $\alpha \in \text{Aut}(A)$ and let $\text{ht}(\alpha)$ denote the Voiculescu's topological entropy of α ([V]). The usual method of computing $\text{ht}(\alpha)$ is based on two steps. First one produces an explicit or semi-explicit approximating net through matrix algebras whose rank can be controlled and thus provides an estimate from above (say given by $M > 0$). Then one looks for an α -invariant C^* -subalgebra $C \subset A$ such that $\text{ht}(\alpha|_C)$ is known to be equal to M . Of course the second step is in a sense recurrent, as we need to be able to determine the entropy of $\alpha|_C$. This becomes much easier when C is a commutative algebra, as then $\alpha|_C$ is induced by a homeomorphism T of the spectrum of C and it was shown in [V] that $\text{ht}(\alpha|_C) = h_{\text{top}}(T)$ – note that the general difficulty in understanding how the positive Voiculescu entropy is produced is reflected in the fact that there is still no direct proof of the inequality $\text{ht}(\alpha_T) \geq h_{\text{top}}(T)$, the argument of [V] exploits the properties of dynamical state entropy and classical variational principle. Other connections between the appearance of a non-zero noncommutative entropy and commutativity can be seen in [HS] (where the occurrence of maximal entropy for a system of subalgebras is related to existence of suitable maximally abelian subalgebras) and in [BDS] (where free shifts are shown to have zero Voiculescu entropy).

One could therefore be tempted to believe that actually

$$(1) \quad \text{ht}(\alpha) = \text{ht}_c(\alpha),$$

where

$$\text{ht}_c(\alpha) = \sup \{ \text{ht}(\alpha|_C) : C - \text{commutative } \alpha - \text{invariant } C^* - \text{subalgebra of } A \}.$$

The aim of this note is to observe that the bit-stream shifts investigated in Chapter 12 of [NS] (see also [GS], [NST]) giving counterexamples to the tensor product formula for Connes-Narnhofer-Thirring entropy also provide counterexamples to the equality in (1).

Before we state the result let us quickly summarise the properties of the bit-stream shifts that we will need in what follows. Let $X \subset \mathbb{N}$. Then X determines in a natural way a bicharacter ω on $G := \bigoplus_{\mathbb{Z}} \mathbb{Z}_2$ and hence a (nuclear) twisted C^* -algebra $A(X) = C^*(G, \omega)$ with a canonical faithful tracial state τ . The algebra $A(X)$ is equipped with a τ -preserving shift-type automorphism σ_X . In Chapter 12

Permanent address of the author: Department of Mathematics, University of Łódź, ul. Banacha 22, 90-238 Łódź, Poland.

2000 *Mathematics Subject Classification.* Primary 46L55.

Key words and phrases. Noncommutative topological entropy, C^* -dynamical systems.

of [NS] it was shown that for almost every (with respect to the Bernoulli measure on $2^{\mathbb{N}} \approx \prod_{n \in \mathbb{N}} \{0, 1\}$) nonperiodic $X \subset \mathbb{N}$ the automorphism σ_X has the following properties:

- (i) τ is the unique σ_X -invariant state on $A(X)$;
- (ii) $h_\tau(\sigma_X) = 0$;
- (iii) $h_{\tau \otimes \tau}(\sigma_X \otimes \sigma_X) = \log 2$.

Here and in what follows $h_\phi(\alpha)$ denotes the noncommutative state entropy of Connes, Narnhofer and Thirring introduced in [CNT] (ϕ is an α -invariant state).

Theorem 1. *For almost every nonperiodic set $X \subset \mathbb{N}$ the binary shift σ_X satisfies $ht_c(\sigma_X) = 0 < \frac{\log 2}{2} \leq ht(\sigma_X)$.*

Proof. Let X be a subset of $\mathbb{N}$ such that σ_X satisfies the conditions (i)–(iii) stated before the theorem. We will write $A := A(X)$, $\sigma := \sigma_X$.

The upper estimate follows immediately from property (iii) above, subadditivity of Voiculescu's topological entropy with respect to tensor products (Theorem 6.2.2 (iv) of [NS]) and the fact that it dominates the CNT entropy (Theorem 6.2.7 in [NS]).

Let then C be a commutative σ -invariant C^* -subalgebra of A . Write σ_C for $\sigma|_C$. Due to the classical variational principle it suffices to show that $h_\phi(\sigma_C) = 0$ for every σ_C -invariant state ϕ on C . We claim first that invariance of ϕ implies it is equal to $\tau|_C$. Indeed, suppose that ψ is an arbitrary state on A extending ϕ . Let $\tilde{\psi}$ be a weak*-limit point of the sequence $(\frac{1}{n} \sum_{i=1}^n \psi \circ \sigma^i)_{n=1}^\infty$. Then $\tilde{\psi}$ is σ -invariant and extends ϕ . By condition (i) above $\tilde{\psi} = \tau$ so $\phi = \tau|_C$.

We are almost done, as due to property (ii) before the statement of the theorem we have $h_\tau(\sigma) = 0$. We need however to remember that the CNT entropy is known to be monotone only under passing to the expected invariant subalgebras and we did not assume that C is expected. The traciality of τ tells us that we have a τ -preserving expectation as soon as we pass to the von Neumann algebraic setting – this will be exploited in the last part of the proof.

Let $(\pi_\tau, L^2(A, \tau), \Omega_\tau)$ be a GNS triple for (A, τ) , let $\tilde{\tau}$ denote the extension of τ to $\pi_\tau(A)''$ (given by the vector state associated with Ω_τ) and let $\tilde{\sigma}$ denote the unique normal automorphism of $\pi_\tau(A)''$ satisfying $\tilde{\sigma}(\pi_\tau(a)) = \pi_\tau(\sigma(a))$ for all $a \in A$. By Theorem 3.2.2 (ii) in [NS]

$$(2) \quad h_\tau(\sigma) = h_{\tilde{\tau}}(\tilde{\sigma}).$$

Note that $\tilde{\sigma}$ leaves $\pi_\tau(C)''$ invariant.

Let $p \in B(L^2(A, \tau))$ be the orthogonal projection onto $K := \text{cl}(\pi_\tau(C)\Omega_\tau)$. Then $p \in \pi_\tau(C)'$. It is easy to see that $(\pi_C := \pi_\tau|_C, K, \Omega_\tau)$ is a GNS triple for $(C, \tau|_C)$. Denote by $\hat{\tau} \in \pi_C(C)''_*$, $\hat{\sigma}_C \in \text{Aut}(\pi_C(C)'')$ corresponding normal extensions of $\tau|_C$ and σ . Again by Theorem 3.2.2 (ii) in [NS]

$$(3) \quad h_{\tau|_C}(\sigma_C) = h_{\hat{\tau}}(\hat{\sigma}_C).$$

It remains to observe that the map $\Psi : \pi_\tau(C)'' \rightarrow \pi_C(C)''$ given by $m \mapsto mp$ is a covariant isomorphism of $\pi_\tau(C)''$ and $\pi_C(C)''$, by which we mean that $\Psi \circ \tilde{\sigma}|_{\pi_\tau(C)''} = \hat{\sigma}_C \circ \Psi$ and $\hat{\tau} \circ \Psi = \tilde{\tau}|_C$. Indeed, the fact that Ψ is injective follows from the faithfulness of τ , whereas the surjectivity and covariance properties are consequences of

the normality of all maps involved and the fact that the corresponding statements clearly hold on $\pi_\tau(\mathbf{C})$. Thus we have

$$(4) \quad h_{\tilde{\tau}|_{\pi_\tau(\mathbf{C})''}}(\tilde{\sigma}|_{\pi_\tau(\mathbf{C})''}) = h_{\tilde{\tau}}(\hat{\sigma}_{\mathbf{C}}).$$

As τ is a trace, there exists a (unique) τ -preserving conditional expectation from $\pi_\tau(\mathbf{A})''$ onto $\pi_\tau(\mathbf{C})''$. Therefore by Theorem 3.2.2 (v) in [NS]

$$(5) \quad h_{\tilde{\tau}|_{\pi_\tau(\mathbf{C})''}}(\tilde{\sigma}|_{\pi_\tau(\mathbf{C})''}) \leq h_{\tilde{\tau}}(\tilde{\sigma}).$$

Putting together equations (2)-(5) yields

$$h_{\tau|_{\mathbf{C}}}(\sigma_{\mathbf{C}}) \leq h_\tau(\sigma) = 0$$

and the proof is finished. □

Note that considering a direct sum of the dynamical system satisfying the conditions in Theorem 1 with a noncommutative dynamical system $(\mathbf{B}, \gamma)$ such that $\text{ht}(\gamma) = \text{ht}_c(\gamma) \in (0, \frac{\log 2}{2})$ one can obtain examples in which $\text{ht}(\alpha) > \text{ht}_c(\alpha) > 0$.

One version of the analogous problem for the Connes-Størmer entropy would be the following – let $\mathbf{M}$ be the hyperfinite II_1 -factor with the trace τ . Does there exist a trace-preserving automorphism $\alpha \in \text{Aut}(\mathbf{M})$ such that

$$h_\tau(\alpha) > \sup \{h_\tau(\alpha|_{\mathbf{C}}) : \mathbf{C} - \alpha - \text{invariant masa in } \mathbf{M}\}?$$

In [GN] the authors gave examples of automorphisms α of the hyperfinite II_1 -factor such that $h_\tau(\alpha) > h_\tau(\alpha|_{\mathbf{C}})$, where $\mathbf{C}$ is a fixed α -invariant Cartan subalgebra of $\mathbf{M}$ (other examples of such situation arise for permutation endomorphisms of Cuntz algebras, as can be deduced from [SZ]), but the problem stated above remains open.

In view of Theorem 1 it is also natural to ask the following question: do there exist a (say unital) C^* -algebra $\mathbf{A}$ and $\alpha \in \text{Aut}(\mathbf{A})$ such that $\text{ht}(\alpha) > 0$ and $\mathbf{A}$ does not admit any nontrivial α -invariant abelian subalgebras?

Acknowledgment. The note was written during a visit of the author to University of Tokyo in October-November 2009 funded by a JSPS Short Term Postdoctoral Fellowship. The author would like to thank Sergey Neshveyev for useful comments on the first draft.

REFERENCES

- [BDS] N. Brown, K. Dykema and D. Shlyakhtenko, Topological entropy of free product automorphisms, *Acta Math.* **189** (2002), 1–35.
- [CNT] A. Connes, H. Narnhofer and W. Thirring, Dynamical entropy of C^* -algebras and von Neumann algebras, *Comm. Math. Phys.* **112** (1987), no. 2, 691–719.
- [GN] V. Ya. Golodets and S. Neshveyev, Entropy of automorphisms of II_1 -factors arising from the dynamical systems theory, *J. Funct. Anal.* **181** (2001), no. 1, 14–28.
- [GS] V. Ya. Golodets and E. Størmer, Entropy of C^* -dynamical systems generated by bitstreams, *Ergodic Theory Dynam. Systems* **18** (1998), 859–874.
- [HS] U. Haagerup and E. Størmer, Maximality of entropy in finite von Neumann algebras, *Invent. Math.* **132** (1998), 433–455.
- [NS] S. Neshveyev and E. Størmer, “Dynamical entropy in operator algebras,” *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics*, 50. Springer-Verlag, Berlin, 2006.
- [NST] H. Narnhofer, E. Størmer and W. Thirring, C^* -dynamical system for which the tensor product formula for entropy fails, *Ergodic Theory Dynam. Systems* **15** (1995), 961–968.

- [SZ] A. Skalski and J. Zacharias, Noncommutative topological entropy of endomorphisms of Cuntz algebras, *Lett. Math. Phys.* **86** (2008), no. 2-3, 115–134.
- [V] D. Voiculescu, Dynamical approximation entropies and topological entropy in operator algebras, *Comm. Math. Phys.* **170** (1995), no. 2, 249–281.

DEPARTMENT OF MATHEMATICS AND STATISTICS, LANCASTER UNIVERSITY, LANCASTER LA1 4YF, UNITED KINGDOM

E-mail address: `a.skalski@lancaster.ac.uk`